



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORBIDITY AND MORTALITY—Continued.

Morbidity and mortality table, cities of the United States, for week ended Mar. 16, 1912—Continued.

Cities.	Popula- tion, United States census, 1910.	Total deaths from all causes. ¹	Diph- theria.		Measles.		Scarlet fever.		Small- pox.		Tuber- culosis.		Ty- phoid fever.	
			Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.	Cases.	Deaths.
<i>Cities having less than 25,000 inhabitants.</i>														
Alameda, Cal.	23,383	5			2									
Beaver Falls, Pa.	12,191		1				1						3	
Bennington, Vt.		4												
Braddock, Pa.	19,957	15	6	2	1									
Cambridge, Ohio	11,327	3										1		
Carbondale, Pa.	17,040	15	1	1	4		2							
Clinton, Mass.	13,075	4												
Coffeyville, Kans.	12,687		2						1					
Columbus, Ga.	20,554	4												
Columbus, Ind.		4					1		3			2		
Concord, N. H.	21,497	8			13									
Cumberland, Md.	21,839	7			1						1	2	1	
Dunkirk, N. Y.		4			1									
Galesburg, Ill.	20,089	7	1						1					
Gloucester, Mass.	24,398	12										1		
Harrison, N. J.	14,498	3	1	1							2			
Kearny, N. J.	18,659	5	1		8		2				2	1		
La Fayette, Ind.	20,081	11			2							1		
Lebanon, Pa.	19,240		2											
Logansport, Ind.	19,050	5			1							1	2	
Marinette, Wis.		4			2								2	
Marlboro, Mass.	14,577	4					1							
Massillon, Ohio		4	2				1							
Medford, Mass.	23,156	6			13		1						1	
Melrose, Mass.	15,715	3			4									
Moline, Ill.	24,199	13										3		
Montclair, N. J.	21,150	8			1		1				1	1		
Morristown, N. J.	12,507	3			3		3				1			
Nanticoke, Pa.	18,507	4			4									
Newburyport, Mass.	19,240		1								2			
North Adams, Mass.	22,012	5									1	1		
Northampton, Mass.	19,431	5									2			
Ottumwa, Iowa.	22,012	11										1		
Palmer, Mass.		2												
Palo Alto, Cal.		1			1									
Plainfield, N. J.	22,050	9			49		1				1		1	
Portsmouth, N. H.			1				1		1					
Pottstown, Pa.		8			1		1							
South Bethlehem, Pa.	19,973				5		1					3		
Warren, Pa.	11,081	4		1										
Wilkinsburg, Pa.	18,924	13	1		2		1							
Woburn, Mass.	15,308	2			12									

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES
OF THE UNITED STATES (Untabulated).

CALIFORNIA.—Month of September, 1911. Population, 2,377,549. Total number of deaths from all causes 2,532, including diphtheria 11, measles 4, scarlet fever 3, tuberculosis 373, typhoid fever 45. Cases reported: Diphtheria 60, measles 232, scarlet fever 69, smallpox 19, tuberculosis (pulmonary), 217, typhoid fever 120.

FLORIDA.—Reports from the State board of health show as follows:

Week ended March 9, 1912. Diphtheria present in 6 localities with 17 cases, malaria in 6 localities with 17 cases, smallpox in 9 counties with 53 cases, tuberculosis in 8 localities with 14 cases, typhoid fever in 4 localities with 10 cases.

Week ended March 16, 1912. Diphtheria present in 4 localities with 12 cases, malaria in 4 localities with 11 cases, smallpox in 9 counties with 33 cases, tuberculosis in 6 localities with 10 cases, typhoid fever in 2 localities with 8 cases.

INDIANA.—Month of January, 1912. Population, 2,700,876. Total number of deaths from all causes 3,186, including diphtheria 24, measles 8, scarlet fever 7, tuberculosis 359, typhoid fever 29. Cases reported: Diphtheria 225 in 41 counties, scarlet fever 294 in 49 counties, smallpox 89 in 15 counties, typhoid fever 131 in 34 counties.

MARYLAND.—Month of February, 1912. Population, 1,295,346. Total number of deaths from communicable diseases 153, including diphtheria 7, scarlet fever 3, tuberculosis 97, typhoid fever 9. Cases reported: Diphtheria 73, measles 100, scarlet fever 64, typhoid fever 57.

MICHIGAN.—Month of February, 1912. Population, 2,810,173. Cases of communicable diseases reported: Diphtheria 199, measles 166, scarlet fever 460, smallpox 95, tuberculosis 147, typhoid fever 203.